

Strong fragmentation of low-energy electromagnetic excitation strength in ^{117}Sn

V. Yu. Ponomarev^{1*}, J. Bryssinck¹, L. Govor², F. Bauwens¹, O. Beck³, D. Belic³, P. von Brentano⁴, D. De Frenne¹, C. Fransen⁴, R.-D. Herzberg^{4,5}, E. Jacobs¹, U. Kneissl³, H. Maser³, A. Nord³, N. Pietralla⁴, H.H. Pitz³, V. Werner⁴

¹*Vakgroep Subatomaire en Stralingsfysica, Universiteit Gent, Proeftuinstraat 86, 9000 Gent, Belgium*

²*Russian Scientific Centre "Kurchatov Institute", Moscow, Russia*

³*Institut für Strahlenphysik, Universität Stuttgart, Stuttgart, Germany*

⁴*Institut für Kernphysik, Universität zu Köln, Köln, Germany.*

⁵*Oliver Lodge Laboratory, University of Liverpool, Oxford Street, Liverpool, L69 7ZE, UK*

(February 8, 2008)

Results of nuclear resonance fluorescence experiments on ^{117}Sn are reported. More than 50 γ transitions with $E_\gamma < 4$ MeV were detected indicating a strong fragmentation of the electromagnetic excitation strength. For the first time microscopic calculations making use of a complete configuration space for low-lying states are performed in heavy odd-mass spherical nuclei. The theoretical predictions are in good agreement with the data. It is concluded that although the $E1$ transitions are the strongest ones also $M1$ and $E2$ decays contribute substantially to the observed spectra. In contrast to the neighboring even $^{116-124}\text{Sn}$, in ^{117}Sn the 1^- component of the two-phonon $[2_1^+ \otimes 3_1^-]$ quintuplet built on top of the $1/2^+$ ground state is proved to be strongly fragmented.

PACS numbers: 21.10.Re, 21.60.-n, 23.20.-g, 25.20.Dc

During the last years the study of the properties of multi-phonon excitations in atomic nuclei has been one of the key topics in nuclear structure research. The main motivation for these studies is to gain insight into the applicability of a picture of harmonic excitations to a finite strongly interacting fermion system, like an atomic nucleus. At present, systematic experimental information on the properties of low-lying two-phonon states [1–3] as well as on two-phonon giant resonances [4,5] in even-even spherical nuclei is available.

The 1^- component of a two-phonon quintuplet built up of collective quadrupole and octupole phonons, $[2_1^+ \otimes 3_1^-]$, is a very good candidate for the investigation of anharmonic properties of low-lying excited states. This state is practically a pure two-phonon state with very weak admixtures of other configurations [3]. In addition, a tool which is very selective to its excitation, namely nuclear resonance fluorescence (NRF), is available. The main conclusion from systematic studies [2,3,6–8] of this 1^- state in even spherical nuclei in different mass regions is that anharmonicity effects are rather weak, supporting a harmonic treatment of nuclear excitations.

The next step in these studies should be the extension to spherical odd-mass nuclei. Here experimental information is sparse. NRF experiments on ^{113}Cd [9], ^{133}Cs [10] and ^{143}Nd [11] have been reported. From very general arguments one expects rather weak changes in the properties of collective vibrations when they are coupled to an odd quasiparticle. However, even the first NRF experiments indicate a rather strong fragmentation of the excitation strength. To minimize the number of possible excited levels it is imperative to consider this coupling of collective vibrations to an odd quasiparticle in a nucleus with ground state spin $1/2$.

In the present letter we report on NRF experiments

on ^{117}Sn ($J_{g.s.}^\pi = 1/2^+$) in which 59 transitions with $E_\gamma < 4.05$ MeV have been observed. It is clear that a detailed theoretical analysis which provides information on the structure of the excited states, is necessary to explain the results of this experiment. For this reason a full microscopic calculation of the properties of multi-phonon states in heavy odd-mass spherical nuclei has been performed for the first time.

Experimental results have been obtained at the bremsstrahlung NRF-facility at the 4.3 MV Dynamitron accelerator of the Stuttgart University [1]. The electron energy was 4.1 MeV. The NRF technique has already been extensively described in review articles (see, e.g. Ref. [1]). For this experiment, a setup with 3 HP Ge-detectors with 100% efficiency (relative to a $3'' \times 3''$ Na(Tl) crystal) and installed at scattering angles 90° , 127° and 150° was used. The NRF target consisted of two tin disks. The total amount of tin was 1.649 g and the enrichment in ^{117}Sn was 92.10%. Two ^{27}Al disks with a total weight of 780 mg were sandwiched between the tin disks. The aluminum nuclei, having well known photon scattering cross sections [12], enable the photon flux calibration and thereby the absolute measurement of the photon scattering cross sections of excited states in the ^{117}Sn nuclei.

In this experiment 59 γ -transitions with an energy between 1440 and 4050 keV have been detected. From the consideration of γ energy differences it follows that only 5 of the observed γ transitions may be due to inelastic photon scattering. The energy integrated cross sections for elastic scattering, I_s , for all observed states are plotted in Fig. 1a. Spin quantum numbers could be assigned to some excited states from angular distribution measurements: for 6 levels the spin value $3/2$ is suggested and for another 5 levels an assignment $1/2(3/2)$, meaning that

1/2 is more likely, is possible. The present NRF setup does not allow to deduce parities for levels in odd-mass nuclei. In the energy region under consideration only a few levels are known from other experiments [13]. More details on the experimental analysis and the complete information on observed transitions will be published in a forthcoming article.




FIG. 1. a) Experimental and b) calculated intensities of γ -decays of low-lying states in ^{117}Sn into the ground state.

Combining the results of the present experimental studies with those for ^{113}Cd [9], ^{133}Cs [10] and ^{143}Nd [11] we conclude that a strong fragmentation of low-energy electromagnetic excitation strength is a general feature in heavy odd-mass nuclei. With the present state-of-the-art of the modern NRF technique alone it is not possible to study in detail the properties of the many levels involved and thus, a theoretical support is needed.

To accomplish this task a theoretical analysis of the properties of low-lying states in ^{117}Sn has been performed within the Quasiparticle-Phonon Model (QPM) [14]. The QPM has been successfully applied to describe the position and the $E1$ excitation probability of the lowest 1^- state in the even-even $^{116-124}\text{Sn}$ isotopes [3]. This state has a two-phonon nature with a contribution of the $[2_1^+ \otimes 3_1^-]_{1^-}$ configuration of 96-99%. A general QPM formalism to treat odd-mass spherical nuclei is presented in review articles [15,16]. A Woods-Saxon potential is used in QPM as an average field for protons and neutrons. Phonons of different multiplicities and parities are obtained by solving RPA equations with a separable form of the residual interaction. The single-particle

spectrum and phonon basis are fixed from calculations in the neighboring even-even nuclear core. No new free parameters are introduced for calculations in the odd-mass nuclei. All matrix elements of the coupling between the different configurations in the wave functions of states in odd-mass nuclei are calculated on a microscopic footing, making use of the internal fermion structure of the phonons and the model Hamiltonian. In the calculations to be presented below we use the same set of parameters as in our previous calculations in ^{116}Sn [3].

In our present calculations the wave functions of the ground state and excited states are mixtures of different “quasiparticle $\otimes N$ -phonon” ($[qp \otimes Nph]$) configurations, where $N = 0, 1, 2, 3$. When $N \geq 1$, qp from different levels of the average field are accounted for. It is only necessary that all configurations have the same total spin and parity. To achieve a correct position of the $[qp \otimes 2ph]$ configurations, in which we are especially interested in these studies, $[qp \otimes 3ph]$ configurations are important. The excitation energies and the contribution of different components from the configuration space to the structure of each excited state are obtained by a diagonalization of the model Hamiltonian on a set of employed wave functions. Interaction matrix elements are calculated to first order perturbation theory. This means that any $[qp \otimes Nph]$ configuration interacts with the $[qp \otimes (N \pm 1)ph]$ ones, but its coupling to $[qp \otimes (N \pm 2)ph]$ configurations is not included in this theoretical treatment. The omitted couplings have non-vanishing interaction matrix elements only in second order perturbation theory. They are much smaller than the ones accounted for and excluded from consideration for technical reasons. An interaction with other $[qp \otimes Nph]$ configurations is taken into account while treating Pauli principle corrections.

The phonon basis includes the phonons with multipolarity and parity $\lambda^\pi = 1^\pm, 2^+, 3^-$ and 4^+ . Several low-energy phonons of each multipolarity are included in the model space. The most important ones are the first collective $2^+, 3^-$ and 4^+ phonons and the ones which form the giant dipole resonance (GDR). Non-collective low-lying phonons of an unnatural parity and natural parity phonons of higher multiplicities are of a marginal importance. To make realistic calculations possible one has to truncate the configuration space. We have done this on the basis of excitation energy arguments. All $[qp \otimes 1ph]$ and $[qp \otimes 2ph]$ with $E_x \leq 6$ MeV, and $[qp \otimes 3ph]$ with $E_x \leq 8$ MeV configurations which do not violate the Pauli principle are included in the model space. The only exception are $[J_{g.s.} \otimes 1^-]$ configurations which have not been truncated at all to treat a core polarization effect due to the coupling of low-energy dipole transitions to the GDR on a microscopic level. Thus, for electric dipole transitions we have not renormalized effective charges and used $e^{\text{eff}}(p) = (N/A)e$ and $e^{\text{eff}}(n) = -(Z/A)e$ values to separate the center of mass motion. For $M1$ transitions we use $g_s^{\text{eff}} = 0.64g_s^{\text{free}}$ as recommended in Ref. [17].

In this all the configurations of importance for the description of low-lying states up to 4 MeV are included in the model space. The dimension of this space depends on the total spin of the excited states, and it varies between 500 and 700.




FIG. 2. Calculated intensities of γ -decays into the ground state in a) ^{116}Sn and b-d) ^{117}Sn . $E2$ decays are plotted by dashed lines in a) and c). $E1$ decays in b) which are predominantly due to $[3s_{1/2} \otimes [2_1^+ \otimes 3_1^-]_{1-}]_{1/2^-, 3/2^-} \rightarrow 3s_{1/2}$ transitions are marked by triangles.

Since only $E1$, $M1$ and $E2$ transitions can be observed in the present experiment only the properties of excited states with $J^\pi = 1/2^\pm$, $3/2^\pm$ and $5/2^+$ will be presented. As the parities of the decaying levels are unknown and the total spin of only a few of them is assigned, the best comparison between theoretical predictions and experimental data should be at the level of the scattering cross sections, I_s . The calculated I_s values:

$$I_s(\lambda) \propto E_x^{2\lambda-1} \cdot B(\lambda) \cdot \frac{\Gamma_{g.s.}}{\Gamma_{tot}}, \quad (1)$$

where Γ are decay widths, are plotted in Fig. 1b. We present here only ground state decays. The inelastic decays are accounted for in a calculation of the total decay width Γ_{tot} . Details concerning the calculations and branching ratios will be presented elsewhere. Supporting the experimental findings our calculations also provide a strong fragmentation of electromagnetic strength. The strongest transitions have $E1$ character, but also $E2$ and

$M1$ excitations yield comparable cross sections. They are presented separately in Fig. 2b,c,d, respectively, in comparison to calculated intensities in the core nucleus, ^{116}Sn (Fig. 2a). The total calculated scattering cross sections, I_s , of the plotted $E1$, $M1$ and $E2$ excitations in Fig 2b-d equal 73, 37 and 42 eVb. The summed cross section of the experimentally observed levels, presented in Fig.1a, equals 137 eVb and agrees within 10% with the theoretically predicted 152 eVb.

The good correspondence between experimental and theoretical results concerning strength fragmentation, position of main groups of levels and total electromagnetic strength make us confident in the theoretical predictions, which will be used below in explaining the main features of the experimentally observed transitions.

Let us first ask whether the unpaired quasiparticle in odd-mass nuclei plays the role of a spectator in determining the properties of low-lying states. To answer this question we compare the structure of the excited states in ^{117}Sn with the structure of the corresponding states in the neighboring even-even core, ^{116}Sn . We observe the most essential changes in the states connected to the ground state by $E1$ transitions. In the even-even core there is only one 1^- configuration with an excitation energy below 4 MeV (line with triangle in Fig. 2a). It has a $[2_1^+ \otimes 3_1^-]_{1-}$ two-phonon nature [3]. This is a general feature in heavy semi-magic even-even nuclei [18]. All other 1^- configurations are more than 1 MeV above. As a result the 1_1^- state has practically pure two-phonon character in semi-magic nuclei. In contrast, the two corresponding configurations in ^{117}Sn $[3s_{1/2} \otimes [2_1^+ \otimes 3_1^-]_{1-}]_{1/2^-, 3/2^-}$ are embedded in other $[qp \otimes 1ph]$ and $[qp \otimes 2ph]$ configurations with the same spin and parity. An interaction with them leads to a strong fragmentation of these two main configurations (see, Fig. 3). All the resulting states are carrying a fraction of the $E1$ excitation strength from the ground state. The main part of the $[3s_{1/2} \otimes [2_1^+ \otimes 3_1^-]_{1-}]_{1/2^-, 3/2^-}$ configurations is concentrated in $3/2^-$ states with an excitation energy of 3.04 MeV (11%), 3.55 MeV (48%) and 3.56 MeV (32%) and in $1/2^-$ states at 3.00 MeV (22%) and 3.63 MeV (63%) (see, lines with triangles in Fig. 2b). The $E1$ strength distribution among low-lying levels is even more complex because $3/2^-$ states at 2.13, 2.33 and 3.93 MeV have a noticeable contribution of the $3p_{3/2}$ one-quasiparticle configuration with a reduced excitation matrix element $\langle 3p_{3/2} || E1 || 3s_{1/2} \rangle$ for which there is no analogue in the even-even core ^{116}Sn . Also the coupling to $[3s_{1/2} \otimes 1_{GDR}^-]$, for treating the core polarization effect, is somewhat different than in the core nucleus, because of a blocking role of the interaction with other configurations (see, also Ref. [19], where only the last type of transitions have been accounted for in the calculations of the $E1$ strength distribution in ^{115}In). The total $B(E1) \uparrow$ strength in the energy region from 2.0 to 4.0 MeV in this calculation

equals $7.2 \cdot 10^{-3} \text{ e}^2\text{fm}^2$. It should be compared to the calculated $B(E1, 0_{g.s.}^+ \rightarrow [2^+ \otimes 3^-]_{1-}) = 8.2 \cdot 10^{-3} \text{ e}^2\text{fm}^2$ in the neighboring ^{116}Sn nucleus [20].




FIG. 3. Contribution, C^2 , of the $[3s_{1/2} \otimes [2_1^+ \otimes 3_1^-]_{1-}]_{J^\pi}$ configuration to wave functions of $J^\pi = 3/2^-$ (solid lines) and $J^\pi = 1/2^-$ (dashed lines) states in ^{117}Sn . The position of the $[2_1^+ \otimes 3_1^-]_{1-}$ state in ^{116}Sn is indicated by arrow.

Positive parity states in ^{117}Sn are deexciting to the $1/2^+$ ground state by $E2$ and $M1$ transitions. The $B(E2) \uparrow$ strength distribution is dominated by the excitation of the $3/2^+$ state at 1.27 MeV and the $5/2^+$ state at 1.49 MeV. The wave functions of these states carry respectively 85% and 60% of the $[3s_{1/2} \otimes 2_1^+]$ configuration. A smaller fraction of this configuration can be found in the $3/2^+$ state at 2.32 MeV (5%) and the $5/2^+$ state at 2.23 MeV (6%). The rather fragmented $E2$ strength at higher energies (Fig. 2c) is mainly due to $[3s_{1/2} \otimes 2_{4,5}^+]$ configurations which are much less collective than the first one. Fragmented $E2$ strength between 2.0 and 4.0 MeV originating from excitation of $2_{4,5}^+$ phonons has been also observed in NRF experiments on the even ^{116}Sn nucleus [20]. It is supported by theoretical calculations (see, dashed lines in Fig. 2a). In the odd-mass ^{117}Sn nucleus it is even more fragmented because of the higher density of configurations. Nevertheless, these $E2$ excitations at high energies contribute appreciably to the reaction cross section, because the $E2$ photon scattering cross section is a cubic function on the excitation energy (see, Eq. (1)).

The $B(M1) \uparrow$ strength in these calculations is concentrated mainly above 3.5 MeV as can be seen in Fig. 2d. The wave functions of all the $1/2^+$ and $3/2^+$ states at these energies are very complex. The main configurations, responsible for the $M1$ strength, are the $[2d_{5/2,3/2} \otimes 2_i^+]$ ones which are excited because of the internal fermion structure of the phonons (similar to $E1 0_{g.s.}^+ \rightarrow [2_1^+ \otimes 3_1^-]_{1-}$ excitations). They have no analogous transitions in even-even nuclei. The configuration $[3s_{1/2} \otimes 1_1^+]$ has an excitation energy of about 4.2 MeV but its contribution to the structure of states below 4 MeV is rather weak. Most of the states with the largest $B(M1) \uparrow$ values have $J^\pi = 1/2^+$.

To conclude, a strong fragmentation of low-energy elec-

tromagnetic excitation strength in ^{117}Sn has been observed in NRF experiments. This observation is understood by microscopic calculations using a complete configurational space for low-lying states. The calculations show that several $E1$, $M1$ and $E2$ excitations lead to comparably large electromagnetic cross sections, which are expected to be observed in this highly sensitive NRF experiment. The structure of decaying states is analyzed, showing that the $[1/2_{g.s.}^+ \otimes [2^+ \otimes 3^-]_{1-}]$ configuration is strongly fragmented. The responsible configuration mixing is calculated to be considerably more complex, than it was suggested previously [11] in order to explain the corresponding fragmentation of electromagnetic excitation strength in the semi-magic nucleus ^{143}Nd .

We thank Prof. K. Heyde for fruitful suggestions to this manuscript. This work is part of the Research program of the Fund for Scientific Research-Flanders. The support by the Deutsche Forschungsgemeinschaft (DFG) under contracts Kn 154-30 and Br 799/9-1 is gratefully acknowledged. V. Yu. P. acknowledges support from the Research Council of the University of Gent.

-
- * Permanent address: Bogoliubov Lab. of Theor. Phys., Joint Inst. for Nucl. Res., 141980, Moscow region, Dubna, Russia, E-mail: vlad@thsun1.jinr.ru
- [1] U. Kneissl, H. H. Pitz, and A. Zilges, *Prog. Part. Nucl. Phys.* **37**, 349 (1996).
 - [2] R.-D. Herzberg *et al.*, *Nucl. Phys.* **A592**, 211 (1995).
 - [3] J. Bryssinck *et al.*, *Phys. Rev. C* **59**, 1930 (1999).
 - [4] T. Aumann, P. F. Bortignon, H. Emling, *Annu. Rev. Nucl. Part. Sci.* **48**, 351 (1998).
 - [5] C. A. Bertulani, V. Yu. Ponomarev, *Phys. Rep.*, in press.
 - [6] M. Wilhelm *et al.*, *Phys. Rev. C* **57**, 577 (1998).
 - [7] J. Enders *et al.*, *Nucl. Phys.* **A636**, 139 (1998).
 - [8] N. Pietralla, *Phys. Rev. C* **59**, 2941 (1999).
 - [9] W. Geiger *et al.*, *Nucl. Phys.* **A580**, 263 (1994).
 - [10] J. Besserer *et al.*, *Phys. Rev. C* **56**, 1276 (1997).
 - [11] A. Zilges *et al.*, *Phys. Rev. Lett.* **70**, 2880 (1993); R.-D. Herzberg *et al.*, *Phys. Rev. C* **51**, 1226 (1995).
 - [12] N. Pietralla *et al.*, *Phys. Rev. C* **51**, 1021 (1995).
 - [13] J. Blachot, G. Marguier, *Nucl. Data Sheets* **66**, 451 (1992).
 - [14] V. G. Soloviev, *Theory of atomic nuclei : Quasiparticles and Phonons* (Institute of Physics Publishing, Bristol and Philadelphia, 1992).
 - [15] A. I. Vdovin *et al.*, *Part. Nucl.* **16**, 245 (1985).
 - [16] S. Gales, Ch. Stoyanov, A. I. Vdovin, *Phys. Rep.* **166**, 125 (1988).
 - [17] P. von Neumann-Cosel *et al.*, *Phys. Rev. Lett.* **82**, 1105 (1999).
 - [18] V. Yu. Ponomarev *et al.*, *Nucl. Phys.* **A635**, 470 (1998).
 - [19] P. von Neumann-Cosel *et al.*, *Z. Phys. A* **350**, 303 (1995).
 - [20] J. Bryssinck *et al.*, APS preprint aps1999apr22-001.